

# B U L L E T I N

DE LA SOCIÉTÉ DES SCIENCES ET DES LETTRES DE ŁÓDŹ  
2018 Vol. LXVIII

---

Recherches sur les déformations

no. 2

pp. 45–52

*Dedicated to the memory of  
Professor Yurii B. Zelinskii*

Anatoly S. Serdyuk and Tetiana A. Stepanyuk

## LEBESGUE-TYPE INEQUALITIES FOR THE FOURIER SUMS ON CLASSES OF GENERALIZED POISSON INTEGRALS

### Summary

For functions from the set of generalized Poisson integrals  $C_{\beta}^{\alpha,r}L_p$ ,  $1 \leq p < \infty$ , we obtain upper estimates for the deviations of Fourier sums in the uniform metric in terms of the best approximations of the generalized derivatives  $f_{\beta}^{\alpha,r}$  of functions of this kind by trigonometric polynomials in the metric of the spaces  $L_p$ . Obtained estimates are asymptotically best possible.

*Keywords and phrases:* Lebesgue-type inequalities, Fourier sums, generalized Poisson integrals, best approximations

Let  $L_p$ ,  $1 \leq p < \infty$ , be the space of  $2\pi$ -periodic functions  $f$  summable to the power  $p$  on  $[0, 2\pi]$ , in which the norm is given by the formula  $\|f\|_p = \left( \int_0^{2\pi} |f(t)|^p dt \right)^{\frac{1}{p}}$ ;  $L_{\infty}$  be the space of measurable and essentially bounded  $2\pi$ -periodic functions  $f$  with the norm  $\|f\|_{\infty} = \text{ess sup}_t |f(t)|$ ;  $C$  be the space of continuous  $2\pi$ -periodic functions  $f$ , in which the norm is specified by the equality  $\|f\|_C = \max_t |f(t)|$ .

Denote by  $C_{\beta}^{\alpha,r}L_p$ ,  $\alpha > 0$ ,  $r > 0$ ,  $\beta \in \mathbb{R}$ ,  $1 \leq p \leq \infty$ , the set of all  $2\pi$ -periodic

functions, representable for all  $x \in \mathbb{R}$  as convolutions of the form (see, e.g., [1, p. 133])

$$f(x) = \frac{a_0}{2} + \frac{1}{\pi} \int_{-\pi}^{\pi} P_{\alpha,r,\beta}(x-t)\varphi(t)dt, \quad a_0 \in \mathbb{R}, \quad \varphi \perp 1, \quad (1)$$

where  $\varphi \in L_p$  and  $P_{\alpha,r,\beta}(t)$  are fixed generated kernels

$$P_{\alpha,r,\beta}(t) = \sum_{k=1}^{\infty} e^{-\alpha k^r} \cos\left(kt - \frac{\beta\pi}{2}\right), \quad \alpha, r > 0, \quad \beta \in \mathbb{R}. \quad (2)$$

The kernels  $P_{\alpha,r,\beta}$  of the form (2) are called generalized Poisson kernels. For  $r = 1$  and  $\beta = 0$  the kernels  $P_{\alpha,r,\beta}$  are usual Poisson kernels of harmonic functions.

If the functions  $f$  and  $\varphi$  are related by the equality (1), then function  $f$  in this equality is called generalized Poisson integral of the function  $\varphi$ . The function  $\varphi$  in equality (1) is called as generalized derivative of the function  $f$  and is denoted by  $f_{\beta}^{\alpha,r}$ .

The set of functions  $f$  from  $C_{\beta}^{\alpha,r}L_p$ ,  $1 \leq p \leq \infty$ , such that  $f_{\beta}^{\alpha,r} \in B_p^0$ , where

$$B_p^0 = \{\varphi : \|\varphi\|_p \leq 1, \varphi \perp 1\}, \quad 1 \leq p \leq \infty,$$

we will denote by  $C_{\beta,p}^{\alpha,r}$ .

Let  $E_n(f)_{L_p}$  be the best approximation of the function  $f \in L_p$  in the metric of space  $L_p$ ,  $1 \leq p \leq \infty$ , by the trigonometric polynomials  $t_{n-1}$  of degree  $n-1$ , i.e.,

$$E_n(f)_{L_p} = \inf_{t_{n-1}} \|f - t_{n-1}\|_{L_p}.$$

Let  $\rho_n(f; x)$  be the following quantity

$$\rho_n(f; x) := f(x) - S_{n-1}(f; x), \quad (3)$$

where  $S_{n-1}(f; \cdot)$  are the partial Fourier sums of order  $n-1$  for a function  $f$ .

Least upper bounds of the quantity  $\|\rho_n(f; \cdot)\|_C$  over the classes  $C_{\beta,p}^{\alpha,r}$ , we denote by  $\mathcal{E}_n(C_{\beta,p}^{\alpha,r})_C$ , i.e.,

$$\mathcal{E}_n(C_{\beta,p}^{\alpha,r})_C = \sup_{f \in C_{\beta,p}^{\alpha,r}} \|f(\cdot) - S_{n-1}(f; \cdot)\|_C, \quad r > 0, \quad \alpha > 0, \quad 1 \leq p \leq \infty. \quad (4)$$

Asymptotic behaviour of the quantities  $\mathcal{E}_n(C_{\beta,p}^{\alpha,r})_C$  of the form (4) was studied in [2]–[10].

In [11]–[13] it was found the analogs of the Lebesgue inequalities for functions  $f \in C_{\beta}^{\alpha,r}L_p$  in the case  $r \in (0, 1)$  and  $p = \infty$ , and also in the case  $r \geq 1$  and  $1 \leq p \leq \infty$ , where the estimates for the deviations  $\|f(\cdot) - S_{n-1}(f; \cdot)\|_C$  are expressed in terms of the best approximations  $E_n(f_{\beta}^{\alpha,r})_{L_p}$ . Namely, in [11] it is proved that the following best possible inequality holds

$$\|f(\cdot) - S_{n-1}(f; \cdot)\|_C \leq \left( \frac{4}{\pi^2} \ln n^{1-r} + O(1) \right) e^{-\alpha n^r} E_n(f_{\beta}^{\alpha,r})_{L_{\infty}}, \quad (5)$$

where  $O(1)$  is a quantity uniformly bounded with respect to  $n$ ,  $\beta$  and  $f \in C_{\beta}^{\alpha,r}L_{\infty}$ .

The present paper is a continuation of [11], [12], and is devoted to getting asymptotically best possible analogs of Lebesgue-type inequalities on the sets  $C_{\beta}^{\alpha,r}L_p$ ,  $r \in (0, 1)$  and  $p \in [1, \infty)$ . This case was not considered yet. Let formulate the results of the paper.

By  $F(a, b; c; d)$  we denote Gauss hypergeometric function

$$F(a, b; c; z) = 1 + \sum_{k=1}^{\infty} \frac{(a)_k(b)_k}{(c)_k} \frac{z^k}{k!},$$

$$(x)_k := \frac{x}{2} \left( \frac{x}{2} + 1 \right) \left( \frac{x}{2} + 2 \right) \dots \left( \frac{x}{2} + k - 1 \right).$$

For arbitrary  $\alpha > 0$ ,  $r \in (0, 1)$  and  $1 \leq p \leq \infty$  we denote by  $n_0 = n_0(\alpha, r, p)$  the smallest integer  $n$  such that

$$\frac{1}{\alpha r} \frac{1}{n^r} + \frac{\alpha r \chi(p)}{n^{1-r}} \leq \begin{cases} \frac{1}{(3\pi)^3} \cdot \frac{p-1}{p}, & p = 1, \\ \frac{1}{(3\pi)^3}, & 1 < p < \infty, \\ \frac{1}{(3\pi)^3}, & p = \infty, \end{cases} \quad (6)$$

where  $\chi(p) = p$  for  $1 \leq p < \infty$  and  $\chi(p) = 1$  for  $p = \infty$ .

The following statement holds.

**Theorem 1.** *Let  $0 < r < 1$ ,  $\alpha > 0$ ,  $\beta \in \mathbb{R}$  and  $n \in \mathbb{N}$ . Then in the case  $1 < p < \infty$  for any function  $f \in C_{\beta}^{\alpha,r}L_p$  and  $n \geq n_0(\alpha, r, p)$ , the following inequality is true:*

$$\begin{aligned} \|f(\cdot) - S_{n-1}(f; \cdot)\|_C &\leq e^{-\alpha n^r} n^{\frac{1-r}{p}} \left( \frac{\|\cos t\|_{p'}}{\pi^{1+\frac{1}{p'}} (\alpha r)^{\frac{1}{p}}} F^{\frac{1}{p'}} \left( \frac{1}{2}, \frac{3-p'}{2}; \frac{3}{2}; 1 \right) + \right. \\ &\quad \left. + \gamma_{n,p} \left( \left( 1 + \frac{(\alpha r)^{\frac{p'-1}{p}}}{p' - 1} \right) \frac{1}{n^{\frac{1-r}{p}}} + \frac{(p)^{\frac{1}{p'}}}{(\alpha r)^{1+\frac{1}{p}}} \frac{1}{n^r} \right) \right) E_n(f_{\beta}^{\alpha,r})_{L_p}, \quad \frac{1}{p} + \frac{1}{p'} = 1, \end{aligned} \quad (7)$$

where  $F(a, b; c; d)$  is Gauss hypergeometric function, and in the case  $p = 1$  for any function  $f \in C_{\beta}^{\alpha,r}L_1$  and  $n \geq n_0(\alpha, r, 1)$ , the following inequality is true:

$$\|f(\cdot) - S_{n-1}(f; \cdot)\|_C \leq e^{-\alpha n^r} n^{1-r} \left( \frac{1}{\pi \alpha r} + \gamma_{n,1} \left( \frac{1}{(\alpha r)^2} \frac{1}{n^r} + \frac{1}{n^{1-r}} \right) \right) E_n(f_{\beta}^{\alpha,r})_{L_1}. \quad (8)$$

In (7) and (8), the quantity  $\gamma_{n,p} = \gamma_{n,p}(\alpha, r, \beta)$  is such that  $|\gamma_{n,p}| \leq (14\pi)^2$ .

*Proof of Theorem 1.* Let  $f \in C_{\beta}^{\alpha,r}L_p$ ,  $1 \leq p \leq \infty$ . Then, at every point  $x \in \mathbb{R}$  the following integral representation is true:

$$\rho_n(f; x) = f(x) - S_{n-1}(f; x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f_{\beta}^{\alpha,r}(t) P_{\alpha,r,\beta}^{(n)}(x-t) dt, \quad (9)$$

where

$$P_{\alpha,r,\beta}^{(n)}(t) := \sum_{k=n}^{\infty} e^{-\alpha k^r} \cos \left( kt - \frac{\beta \pi}{2} \right), \quad 0 < r < 1, \quad \alpha > 0, \quad \beta \in \mathbb{R}. \quad (10)$$

The function  $P_{\alpha,r,\beta}^{(n)}(t)$  is orthogonal to any trigonometric polynomial  $t_{n-1}$  of degree not greater than  $n-1$ . Hence, for any polynomial  $t_{n-1}$  from we obtain

$$\rho_n(f; x) = \frac{1}{\pi} \int_{-\pi}^{\pi} \delta_n(t) P_{\alpha,r,\beta}^{(n)}(x-t) dt, \quad (11)$$

where

$$\delta_n(x) = \delta_n(\alpha, r, \beta, n; x) := f_{\beta}^{\alpha,r}(x) - t_{n-1}(x). \quad (12)$$

Further we choose the polynomial  $t_{n-1}^*$  of the best approximation of the function  $f_{\beta}^{\alpha,r}$  in the space  $L_p$ , i.e., such that

$$\|f_{\beta}^{\alpha,r} - t_{n-1}^*\|_p = E_n(f_{\beta}^{\alpha,r})_{L_p}, \quad 1 \leq p \leq \infty,$$

to play role of  $t_{n-1}$  in (11). Thus, by using the inequality

$$\left\| \int_{-\pi}^{\pi} K(t-u) \varphi(u) du \right\|_C \leq \|K\|_{p'} \|\varphi\|_p, \quad (13)$$

$$\varphi \in L_p, \quad K \in L_{p'}, \quad 1 \leq p \leq \infty, \quad \frac{1}{p} + \frac{1}{p'} = 1$$

(see, e.g., [14, p. 43]), we get

$$\|f(\cdot) - S_{n-1}(f; \cdot)\|_C \leq \frac{1}{\pi} \|P_{\alpha,r,\beta}^{(n)}\|_{p'} E_n(f_{\beta}^{\alpha,r})_{L_p}. \quad (14)$$

For arbitrary  $v > 0$  and  $1 \leq s \leq \infty$  assume

$$\mathcal{I}_s(v) := \left\| \frac{1}{\sqrt{t^2 + 1}} \right\|_{L_s[0,v]}, \quad (15)$$

where

$$\|f\|_{L_s[a,b]} = \begin{cases} \left( \int_a^b |f(t)|^s dt \right)^{\frac{1}{s}}, & 1 \leq s < \infty, \\ \text{ess sup}_{t \in [a,b]} |f(t)|, & s = \infty. \end{cases}$$

It follows from the paper [9] for arbitrary  $r \in (0, 1)$ ,  $\alpha > 0$ ,  $\beta \in \mathbb{R}$ ,  $1 \leq s \leq \infty$ ,  $\frac{1}{s} + \frac{1}{s'} = 1$ ,  $n \in \mathbb{N}$  and  $n \geq n_0(\alpha, r, s')$  the following estimate holds

$$\begin{aligned} \frac{1}{\pi} \|P_{\alpha,r,\beta}^{(n)}\|_s &= e^{-\alpha n^r} n^{\frac{1-r}{s'}} \left( \frac{\|\cos t\|_s}{\pi^{1+\frac{1}{s'}} (\alpha r)^{\frac{1}{s'}}} \mathcal{I}_s \left( \frac{\pi n^{1-r}}{\alpha r} \right) + \right. \\ &\quad \left. + \delta_{n,s}^{(1)} \left( \frac{1}{(\alpha r)^{1+\frac{1}{s'}}} \mathcal{I}_s \left( \frac{\pi n^{1-r}}{\alpha r} \right) \frac{1}{n^r} + \frac{1}{n^{\frac{1-r}{s'}}} \right) \right), \end{aligned} \quad (16)$$

where the quantity  $\delta_{n,s}^{(1)} = \delta_{n,s}^{(1)}(\alpha, r, \beta)$ , satisfies the inequality  $|\delta_{n,s}^{(1)}| \leq (14\pi)^2$ .

Substituting  $s = p' = \infty$ , from 14 and (16) we get (8).

Further, according to [9] for  $n \geq n_0(\alpha, r, s')$ ,  $1 < s < \infty$ ,  $\frac{1}{s} + \frac{1}{s'} = 1$ , the following equality takes place

$$\mathcal{I}_s\left(\frac{\pi n^{1-r}}{\alpha r}\right) = F^{\frac{1}{s}}\left(\frac{1}{2}, \frac{3-s}{2}; \frac{3}{2}; 1\right) + \frac{\Theta_{\alpha,r,s',n}^{(1)}}{s-1} \left(\frac{\alpha r}{\pi n^{1-r}}\right)^{s-1}, \quad (17)$$

where  $|\Theta_{\alpha,r,s',n}^{(1)}| < 2$ .

Let now consider the case  $1 < p < \infty$ .

Formulas (16) and (17) for  $s = p'$  and  $n \geq n_0(\alpha, r, p)$  imply

$$\begin{aligned} \frac{1}{\pi} \|\mathcal{P}_{\alpha,r,n}\|_{p'} &= e^{-\alpha n^r} n^{\frac{1-r}{p}} \left( \frac{\|\cos t\|_{p'}}{\pi^{1+\frac{1}{p'}} (\alpha r)^{\frac{1}{p}}} F^{\frac{1}{p'}}\left(\frac{1}{2}, \frac{3-p'}{2}; \frac{3}{2}; 1\right) + \right. \\ &\quad \left. + \gamma_{n,p}^{(1)} \left( \frac{1}{p'-1} \frac{(\alpha r)^{\frac{p'-1}{p}}}{n^{(1-r)(p'-1)}} + \frac{p^{\frac{1}{p'}}}{(\alpha r)^{1+\frac{1}{p}}} \frac{1}{n^r} + \frac{1}{n^{\frac{1-r}{p}}} \right) \right) = \\ &= e^{-\alpha n^r} n^{\frac{1-r}{p}} \left( \frac{\|\cos t\|_{p'}}{\pi^{1+\frac{1}{p'}} (\alpha r)^{\frac{1}{p}}} F^{\frac{1}{p'}}\left(\frac{1}{2}, \frac{3-p'}{2}; \frac{3}{2}; 1\right) + \right. \\ &\quad \left. + \gamma_{n,p}^{(2)} \left( \left(1 + \frac{(\alpha r)^{\frac{p'-1}{p}}}{p'-1}\right) \frac{1}{n^{(1-r)(p'-1)}} + \frac{p^{\frac{1}{p'}}}{(\alpha r)^{1+\frac{1}{p}}} \frac{1}{n^r} \right) \right), \end{aligned} \quad (18)$$

where the quantities  $\delta_{n,p}^{(i)} = \delta_{n,p}^{(i)}(\alpha, r, \beta)$ , satisfy the inequality  $|\delta_{n,p}^{(i)}| \leq (14\pi)^2$ ,  $i = 1, 2$ . Estimate (7) follows from (14) and (18). Theorem 1 is proved.  $\square$

It should be noticed, that estimates (7) and (8) are asymptotically best possible on the classes  $C_{\beta,p}^{\alpha,r}$ ,  $1 \leq p < \infty$ .

If  $f \in C_{\beta,p}^{\alpha,r}$ , then  $\|f_\beta^{\alpha,r}\|_p \leq 1$ , and  $E_n(f_\beta^{\alpha,r})_{L_p} \leq 1$ ,  $1 \leq p < \infty$ . Considering the least upper bounds of both sides of inequality (7) over the classes  $C_{\beta,p}^{\alpha,r}$ ,  $1 < p < \infty$ , we arrive at the inequality

$$\begin{aligned} \mathcal{E}_n(C_{\beta,p}^{\alpha,r})_C &\leq e^{-\alpha n^r} n^{\frac{1-r}{p}} \left( \frac{\|\cos t\|_{p'}}{\pi^{1+\frac{1}{p'}} (\alpha r)^{\frac{1}{p}}} F^{\frac{1}{p'}}\left(\frac{1}{2}, \frac{3-p'}{2}; \frac{3}{2}; 1\right) + \right. \\ &\quad \left. + \gamma_{n,p} \left( \left(1 + \frac{(\alpha r)^{\frac{p'-1}{p}}}{p'-1}\right) \frac{1}{n^{\frac{1-r}{p}}} + \frac{(p)^{\frac{1}{p'}}}{(\alpha r)^{1+\frac{1}{p}}} \frac{1}{n^r} \right) \right) E_n(f_\beta^{\alpha,r})_{L_p}, \quad \frac{1}{p} + \frac{1}{p'} = 1. \end{aligned} \quad (19)$$

Comparing this relation with the estimate of Theorem 4 from [9] (see also [10]), we conclude that inequality (7) on the classes  $C_{\beta,p}^{\alpha,r}$ ,  $1 < p < \infty$ , is asymptotically best possible.

In the same way, the asymptotical sharpness of the estimate (8) on the class  $C_{\beta,1}^{\alpha,r}$  follows from comparing inequality

$$\mathcal{E}_n(C_{\beta,p}^{\alpha,r})_C \leq e^{-\alpha n^r} n^{1-r} \left( \frac{1}{\pi \alpha r} + \gamma_{n,1} \left( \frac{1}{(\alpha r)^2} \frac{1}{n^r} + \frac{1}{n^{1-r}} \right) \right) E_n(f_\beta^{\alpha,r})_{L_1} \quad (20)$$

and formula (9) from [8].

## Acknowledgements

Second author is supported by the Austrian Science Fund FWF project F5503 (part of the Special Research Program (SFB) Quasi-Monte Carlo Methods: Theory and Applications)

## References

- [1] A. I. Stepanets, Methods of Approximation Theory. VSP: Leiden, Boston, 2005.
- [2] A. I. Stepanets, Deviation of Fourier sums on classes of infinitely differentiable functions, Ukr. Mat. J. **36**:6 (1984), 567–573.
- [3] S. M. Nikolskii, Approximation of functions in the mean by trigonometrical polynomials, (in Russian) Izv. Akad. Nauk SSSR, Ser. Mat. **10** (1946), 207–256.
- [4] S. B. Stechkin, An estimate of the remainder term of Fourier series for differentiable functions, (in Russian) Tr. Mat. Inst. Steklova **145** (1980) 126–151.
- [5] S. A. Telyakovskii, Approximation of functions of high smoothness by Fourier sums, Ukr. Math. J. **41**:4 (1989), 444–451.
- [6] A. I. Stepanets, Deviations of Fourier sums on classes of entire functions, Ukr. Math. J. **41**:6 (1989), 672–677.
- [7] A. S. Serdyuk, Approximation of classes of analytic functions by Fourier sums in the uniform metric, Ukr. Math. J. **57**:8 (2005), 1079–1096.
- [8] A. S. Serdyuk and T. A. Stepanyuk, Uniform approximations by Fourier sums on the classes of convolutions with generalized Poisson kernels, Dop. Nats. Akad. Nauk. Ukr. No. 11 (2016), 10–16.
- [9] A. S. Serdyuk, T. A. Stepanyuk, Approximations by Fourier sums of classes of generalized Poisson integrals in metrics of spaces  $L_s$ , Ukr. Mat. J. **69**:5 (2017), 811–822.
- [10] A. S. Serdyuk, T. A. Stepanyuk, Uniform approximations by Fourier sums on classes of generalized Poisson integrals, Analysis Mathematica, 2018 (accepted for publication)
- [11] A. I. Stepanets, On the Lebesgue inequality on classes of  $(\psi, \beta)$ -differentiable functions, Ukr. Math. J. **41**:4 (1989), 435–443.
- [12] A. I. Stepanets, A. S. Serdyuk, Lebesgue inequalities for Poisson integrals, Ukr. Math. J. **52**:6 (2000), 798–808.
- [13] A. S. Serdyuk, A. P. Musienko, The Lebesgue type inequalities for the de la Vall?e Poussin sums in approximation of Poisson integrals, Zb. Pr. Inst. Mat. NAN Ukr. **7**:1 (2010), 298–316.
- [14] N. P. Korneichuk, Exact Constants in Approximation Theory, Vol. **38**, Cambridge Univ. Press, Cambridge, New York 1990.

Institute of Mathematics  
National Academy of Sciences of Ukraine  
Tereshchenkivska str. 3, UA-01004, Kyiv  
Ukraine  
E-mail: serdyuk@imath.kiev.ua

Presented by Adam Paszkiewicz at the Session of the Mathematical-Physical Commission of the Łódź Society of Sciences and Arts on May 14, 2018.

## NIERÓWNOŚĆ TYPU LEBESGUE'A DLA SUM FOURIERA NA KLASACH UOGÓLNIONYCH CAŁEK POISSONA

### S t r e s z c z e n i e

Dla funkcji ze zbioru uogólnionych całek Poissona  $C_{\beta}^{\alpha,r}L_p$ ,  $1 \leq p < \infty$ , otrzymujemy górne oszacownie dla odchyleń sum Fouriera w jednostajnej metryce w terminach najlepszej aproksymacji uogólnionych pochodnych  $f_{\beta}^{\alpha,r}$  funkcji tego typu w metryce przestrzeni  $L_p$ . Uzyskane oszacowania są asymptotycznie najlepsze z możliwych.

*Slowa kluczowe:* nierówności typu Lebesgue'a, sumy Fouriera, uogólnione całki Poissona, najlepsze przybliżenia

